



Figure 2. EtCO₂/paCO₂ ratio variations during the pronation and one hour after supination in the first and in the second prono-supination maneuver.

Conclusions

The little amount of data limits the statistical significance of our study; nevertheless, our work gives new insights on the suitability of repeated pronation maneuvers. Although simple, the pronation is not free of complications and requires an additional burden in an over-ceiling work environment, such as during the COVID-19 pandemic.

References

- Gattinoni L, Busana M, Giosa L, et al. Prone positioning in Acute Respiratory Distress Syndrome. *Semin Respir Crit Care Med* 2019;40:94-100.
- Abroug F, Ouannes-Besbes L, Elatrous S, et al. The effect of prone positioning in acute respiratory distress syndrome or acute lung injury: a meta-analysis: areas of uncertainty and recommendations for research. *Intensive Care Med* 2008;34:1002-11.
- Sud S, Friedrich JO, Adhikari NK, et al. Effect of prone positioning during mechanical ventilation on mortality among patients with acute respiratory distress syndrome: a systematic review and meta-analysis. *CMAJ* 2014;186:E381-E390.
- Guérin C, Reignier J, Richard JC, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med* 2013;368:2159-68.
- Petrucci N, De Feo C. Lung protective ventilation strategy for the acute respiratory distress syndrome. *Cochrane Database Syst Rev* 2013;2:CD003844.
- Brochard L, Slutsky A, Pesenti A. Mechanical ventilation to minimize progression of lung injury in acute respiratory failure. *Am J Respir Crit Care Med* 2017;195:438-442.
- Qadri SK, Ng P, Toh TSW, et al. Critically ill patients with COVID-19: a narrative review on prone position. *Pulm Ther* 2020;6:233-46.
- Scaramuzza G, Gamberini L, Tonetti T, et al. Sustained oxygenation improvement after first prone positioning is associated with liberation from mechanical ventilation and mortality in critically ill COVID-19 patients: a cohort study. *Ann Intensive Care* 2021;11:63.
- Bussani R, Schneider E, Zentilin L. Persistence of viral RNA, pneumocyte syncytia and thrombosis are hallmarks of advanced COVID-19 pathology. *E Bio Medicine* 2020;61:103-4.
- Gattinoni L, Chiumello D, Rossi S. COVID-19 pneumonia: ARDS or not? *Crit Care* 2020;24:154.
- Harris DE, Massie M. Role of alveolar-arterial gradient in partial pressure of oxygen and PaO₂/fraction of inspired oxygen ratio measurements in assessment of pulmonary dysfunction. *AANA J* 2019;87:214-21.